

# Effectiveness Dose of injection acetaminophen for Control of Shivering after general Anesthesia in inguinal hernia Surgery

Mohammad Sadra Nazari<sup>1</sup>, Asghar karbord<sup>2\*</sup>, Ali Reza Mohammad zade<sup>3</sup>

<sup>1</sup>Assistant Professor of general surgery and vascular surgery, Faculty Member, Department of general surgery, School of Medicine, Qazvin University of Medical Sciences.Iran

<sup>2</sup>Epidemiologist and Faculty Member of Department of Surgical Technologist, School of Paramedical Sciences, Qazvin University of Medical Sciences.Iran

<sup>3</sup>Assistant Professor of general surgery and thorax surgery, Faculty Member, Department of general surgery, School of Medicine, Qazvin University of Medical Sciences.Iran

**\*Corresponding Author:** Asghar karbord. Epidemiologist and Faculty Member of the Department of Surgical Technologist, School of Paramedical Sciences, Qazvin University of Medical Sciences. Iran.

**Received Date:** 30 December 2022; **Accepted Date:** 07 January 2023; **Published date:** 18 January 2023

**Citation:** Asghar karbord, Mohammad Sadra Nazari, Ali Reza Mohammad zade, (2023). Effectiveness Dose of injection acetaminophen for Control of Shivering after general Anesthesia in inguinal hernia Surgery, Journal of Internal Medicine & Health Affairs. 2(1). DOI: 10.58489/2836-2411/011

**Copyright:** © 2023 Asghar karbord, this is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

Inguinal hernia is the most common type of abdominal hernia which is 7 times more common in men than women. Surgical treatment is performed under general anesthesia. about 40 to 60 percent of patients of post-surgery are experiencing tremor. these patients after anesthesia, shivering is very undesirable and from the physiological aspect is very stressful, mild tremor increases the oxygen consumption like a light physical activity and judder increases the metabolic rate and oxygen consumption up to 600%. injection acetaminophen is a safe drug, in recent study determines of Effectiveness dose for control of shivering after anesthesia in surgeries. The aim of this study was to determine the effect of injection acetaminophen on the quality and timing of the Shivering after on the inguinal herniation surgery. This is a randomized double-blind study, that the statistical population was patients that were candidates for general surgery, that were referred to hospitals of Qazvin University of Medical Sciences for treatment. The patient's undergone surgery with general anesthesia with number 1 and 2 anesthetic Class based on anesthesiologist detection. From 80 patients that were randomly selected, 40 people were selected for experimenting group with injection acetaminophen with IM injection of 0.5 mg/kg and 40 people were selected for the placebo control group (routine order). The shivering in the experimenting group was decreased than the control group and was statistically significant. Tremor intensity in the experimental group was lower than the control group in all three measures (1, 2) which was also significant for comparing the linear regression, relative risk (RR) was 8 in two groups according to the normal distribution of the data, in the age group between 20 to 40 years the minimum range of tremor was observed and in the age group that were younger than 20 years range of tremor was observed. Injection acetaminophen (IM) after general anesthesia can reduce the amount and severity of shivering at post- surgery and can be an effective choice drug in the post-surgery in patients with inguinal herniation surgery.

**Keywords:** Inguinal herniation, injection acetaminophen and general anesthesia

## Introduction

An inguinal hernia is a protrusion of the contents of the abdomen in the area of the inguinal canal. The main cause of abdominal hernias is a decrease or weakness in the abdominal wall or increased pressure on it in one area. Inguinal hernia is the most common type of abdominal hernia, which is 7 times more common in men than women. Inguinal hernia

repair is one of the most common surgeries performed by general surgeons and its repair can be performed under general, typical (spinal) anesthesia or local anesthesia [1]. Considering that spinal anesthesia is a safe method of anesthesia for surgery, but about 40 to 60 percent of patients are experiencing shivering. for patients undergoing new methods of spinal anesthesia, shivering is very undesirable and from the physiological aspect is very

stressful, mild tremor increases the oxygen consumption like a light physical activity and judder increases the metabolic rate and oxygen consumption up to be 600%. This case causes a reduction in the amount of arterial oxygen, metabolic acidosis, increased intraocular pressure and increased intracranial pressure, and ECG monitoring, pulse rate, blood pressure and pulseoximetry are interference track in this situation [2]. A variety of methods exist to control shivering during spinal anesthesia: Such as non-pharmacological methods that are devices and equipments that are used to maintain body temperature significantly, but this equipment may be expensive and not practical in all situations. In the pharmacological methods, several drugs have been studied for the prevention or treatment of tremor, including pethidine, flax serine, sufentanil, alfentanil, tramadol, Nefopam, physostigmine, Doksapram, clonidine and Nalbofen. These days the attempt is to use the drugs that are simple cost-effective and readily available. Although the mechanism of all drugs is not fully understood but the impact on body temperature regulation center, opiate receptors, antagonists' control of N-methyl-D-aspartate receptor in the regulation of body temperature reduces the tremor and shivering. acetaminophen is a safe drug, which is cheap and available that can inhibit shivering by controlling the competitive antagonist of -methyl-D-aspartate receptor. In recent studies the effects of acetaminophen alone or with other medications to reduce tremor during and after surgery with different doses (from 0.5 mg/ kg) has been approved but the doses, have side effects such as hallucinations, that appropriate dosage in reducing tremor needs to be obtained by appropriate research. Other studies compared the effect of acetaminophen, pethidine and tramadol with comfort, tremor and temperature inside the esophagus that showed the impact of three drugs to prevent tremor during spinal anesthesia, but better hemodynamic stability and having less side effects of acetaminophen, is superior to the two other drugs. In a study in 2013 in South Africa on 90 patients with 60-18 years that anesthesia for lower limb surgery was performed, it was found that the effect of low-dose midazolam (0.2 mg/kg) with acetaminophen in prevention of tremor is equal to the effect of higher doses of midazolam by 0.25 mg/kg with acetaminophen. The effect of acetaminophen in cesarean section with spinal anesthesia revealed that 0.4 mg/kg acetaminophen, as much as 0.6 mg/kg acetaminophen is effective in preventing tremor. In Iran a study in the field of contrasting the effect of different doses of acetaminophen and pethidine in

reducing the tremor after cesarean surgery was performed, the results showed that: Although acetaminophen significantly controls the tremor but pethidine is still a better choice for this complication. Therefore, in this study the effect of intramuscular acetaminophen at a dose of 0.5 mg/kg to prevent tremor in patients undergoing spinal anesthesia were evaluated [3-16].

### Materials and Methods

This is a randomized double-blind study, that the statistical population was patients that were candidates for inguinal hernia Surgery with the age group between 20 and 60 years that were referred to hospitals of Qazvin University of Medical Sciences for treatment. Patients with emergency surgery, receiving blood and blood products, surgery less than half an hour and more than two hours, further period of anesthesia during surgery that requires medical intervention (drug) and those that have a history of chronic disease and drug consumption were excluded. After obtaining permission from the ethics committee of the University and necessary information, the consent of the patients has completed and the patient's undergone surgery with general anesthesia with number [1,2] and 3 anesthetic Class based on anesthesiologist detection. From 80 patients that were randomly selected, 40 patients were selected for experimenting group with acetaminophen IM injection related to Rotex of Germany and 40 patients were selected for the placebo control group (routine order). The surgery with general anesthesia was performed. Then the patient post-surgery was observed with the help of anesthesiologist using a visual scale of shivering or not and the degree of tremor

(From zero to three), and required data were recorded in the check list. The tremor degree was determined as followed: - grade number zero was without shivering -grade number one was classy, mild tremor of the face and neck - grade number two was visible tremor in more than one muscle group - grade number three was the whole-body muscle activity [14]. Then the data were given to the SPSS29 software and was analyzed with independent t-test and regression. This plan was approved by the Ethics Committee in the Vice Chancellor for Research of Qazvin University of Medical Sciences and Health Services with the ethics code IR.QUMS.REC.1399.112 on 8/12/99.

Results

Table 1: Shivering category in patients

Shivering degree	Clinical signs and symptoms of shivering
0	Without shivering
1	Low-grade shivering of face and neck
2	Visible shivering in more than one muscle
3	Intense muscular activity in whole body

Table 2: The degree of shivering in both control group and experimenting group

group shivering	frequency	0 degree of tremor	1 degree of tremor	2 degrees of tremor	3 degrees of tremor	Total	p-value
Experimenting group	Number of patients	19	14	6	1	40	0.003
	Percentage of patients	47%	35%	15%	2.5%	100%	
Control group	Number of patients	7	17	11	5	40	
	Percentage of patients	17.5	42.5%	27.5%	12.5%	100%	
Total	number	26	31	17	6	80	
	percent	32.5%	38.75%	21%	7.5%	100%	

Table 3: The relation of dynamic variables with the degree of shivering in studied group

Shivering Group/variable	Group	Without shivering		With shivering (from 1-3 degrees)		p-value	
		Number	Percentage	Number	Percentage		
Experimenting group	sex	men	11	58%	8	38%	0.04
		women	8	42%	13	62%	
	Time of surgery	≤1hour	13	68.5%	5	24%	0.001
		>1hour	6	31.5%	16	76%	
	age	≤20y/o	5	26%	8	38%	0.02
		20-40	11	58%	4	19%	
Control group	sex	40-60	3	15%	9	43%	0.036
		women	5	72%	20	61%	
	Time of surgery	men	2	28%	13	39%	0.0012
		women	5	72%	20	61%	
	age	≤1hour	3	43%	10	30%	0.0012
		>1hour	4	57%	23	70%	
age	≤20y/o	2	28%	6	18%	0.73	
	20-40	2	28%	20	60%		
	40-60	3	43%	7	21%		

### Discussion and Conclusion

In this study, 40 patients were selected for experimenting group and 40 patients were selected for the control group, then comparing the two groups according to the normal distribution of the data, t-test independent and post hoc test were performed. The mean age in the control group was  $[30 \pm 77]$  y/o and the mean age in the intervention group was  $[29 \pm 47]$  y/o. A significant difference with the age was observed ( $p = 0.02$ ), in the age group between 20 to 40 years the minimum range of tremor was observed and in the age group that were younger than 20 years and older than 56 years, the maximum range of tremor was observed. Most people were in the age group of 20-40 years (60%) with time surgery more than 1hour. The shivering severity of the patient's body was recorded in the anesthesia class (class one, two, three) during the surgery by an anesthesiologist, according to the physiological conditions and (Table 1). In this study, all hemodynamic parameters were similar for both groups, except IM injection acetaminophen; anesthesia and the surgical procedures were performed in the operating room under the same conditions. The severity and timing Shivering in the experimenting group was lower than the control group and was statistically significant ( $p = 0.003$ ). Shivering intensity in the experimental group was lower than the control group in all three measures [1, 2 and 3], which was also significant ( $p = 0.003$ ). According to (Table 2), from 40patients in the control group 33% of patients had no chills and 77% experienced the chill. From 40 patients in the experimenting group, 47% had no chills and 53% experienced the chill. With injection acetaminophen of 0.5mg/ kg, reducing from grade number 3 to zero was tangible ( $P = 0.001$ ). The results of this study matched with the study of Mr. Bhatnagar and colleagues confirm the effects of injection acetaminophen doses in reducing the chill during spinal surgery. Also in this study, according to the (Table 3), significant correlation between patient's ages and reduction in chills with injection acetaminophen in both groups, were observed. In the experimental group at an early age, the intensity of shivering increased but then it was considerably decreased and again at 40 years of age, an increase in shivering was observed. Which has been confirmed in other clinical trials [12-17] According to the table (3), in this study, using regression analysis, significant correlation was observed between shivering in both groups and time of surgery ( $P = 0.0012$ ), the highest chills were in more than 1hour time of surgery less than 1 hour time of surgery and the lowest chills was in patient. The tremor was also

higher in women than men, and this co-relation was statistically significant ( $P = 0.02$ ). This relationship shows that women are more prone to chills and necessary injection acetaminophen is more effective in this group. Because of low price and availability of injection acetaminophen, this drug is distinguished in this context. But the determination of a single dose of injection acetaminophen according to hallucinogenic effects of this drug after the patient's [www.opastonline.com](http://www.opastonline.com) J Anesth Pain Med, 2019 Volume 4 | Issue 4 | 3 of 5 recovery causes extensive research on the drug's dose. In this study, it was shown that the effect of injection acetaminophen (0.5mg/ kg) compared to the standard that was set for other studies (0.3- 0.6. mg/kg) was effective in reducing the intensity of chills during the surgery with general anesthesia and controlling factors such as gender and type of surgery and anesthesia class can be very effective (22). Also in this study, a comparison between the types of anesthetic drugs and injection acetaminophen dose with shivering was not performed but according to previous studies (24). This factor can be very effective that by choosing a suitable auxiliary drug in spinal anesthesia (e.g., low dose of petedin or diazepam) that does not have a synergic or antagonistic interfere with the injection acetaminophen, shivering during the surgery can be controlled [20-25]. Hemodynamic status such as temperature, oxygen consumption, cell metabolism and electrolytic stability can be maintained with controlling the chill. The effective injection acetaminophen with the amount of 0.5 mg per kg of body weight to control or reduce the severity of shivering in this study had the same preventing effect in reducing the chill in patients who had undergone the general anesthesia with the study that was conducted by Coase and colleagues, with the injection acetaminophen of 0.6 mg per kg of body weight up to 0.5 mg per kilogram of body. This study and other studies in addition to demonstrating the efficacy of acetaminophen in reducing the shivering, which confirms the results of our study, emphasized on achieving a single and effective dose [13-18]. The relationship between age and the dose of injection acetaminophen in both groups with decreasing the chill and its severity was observed that the amount of shivering in the age groups under 20 and more than 60 years was more than the age groups between 20 and 60 years. In other studies, such as Miller and colleagues it was found out that two factors have played a decisive role in the incidence of tremor: 1 - Young people 2 – reduction in central body temperature. Also, it is believed that with the increasing in the age of patients, the threshold of

shivering, which in normal condition is at a temperature of 36 ° C, decreases one ° C and reaches to 35 ° C. In other words, shivering is higher in the old and adolescents age group. injection acetaminophen was effective for both the upper and lower age limit that corresponds with the results of the study. Considering the prevalence of shivering among adolescents and old people, other methods of analgesia or anesthesia should be used for this age group or the necessary conditions (operating room temperature regulation function and intravenous fluids and blood, and the use of cover heating) for tremor controlling during spinal anesthesia for the patient - before anesthesia –should be provided. Other notable results from this study were the association of gender with shivering (P = 0.06), which is in same direction with available resources. In both groups, as well as control group, shivering in women was more than men. Reduction and control of shivering during and after surgery with general anesthesia should be determined with controlling the effective factors such as the type of anesthesia drug, age, gender and type of surgery in the study. injection acetaminophen at a dose of 0.5mg/kg in after general anesthesia can reduce the amount of shivering after the surgery and can be an effective choice in the field of inguinal hernia surgery for candidates over 60 years of age or in women who are candidate for surgery.

## References

1. Fitzgibbons RJ, Forse, RA . Clinical practice. Groin hernias in adults. *The New England Journal of Medicine*.2015. 372 (8): 756–63.
2. Karbord, A., Kayalha, H., Roushanfekr, M. Q., & Mehdipoor, H. (2019). Effectiveness Dose of Ketamine for Control of Shivering During Spinal Anesthesia in Surgeries. *Journal of Anesthesia & Pain Medicine*, 4(4), 1-5.
3. De Witte, J., & Sessler, D. I. (2002). Perioperative shivering: physiology and pharmacology. *The Journal of the American Society of Anesthesiologists*, 96(2), 467-484.
4. Bhatnagar, S., Saxena, A., Kannan, T. R., Punj, J., Panigrahi, M., & Mishra, S. (2001). Tramadol for postoperative shivering: a double-blind comparison with pethidine. *Anaesthesia and intensive care*, 29(2), 149-154.
5. Katyal Sunil, Tewari Anurag (2002) Shivering: Anesthesia considerations. *J Anesthesia Clinical pharmacology*. 2002; 18: 363-376.
6. Mathews, S., Al Mulla, A., Varghese, P. K., Radim, K., & Mumtaz, S. (2002). Postanaesthetic shivering-a new look at tramadol. *Anaesthesia*, 57(4), 387-403.
7. Sagir, O., Gulhas, N. U. R. Ç. İ. N., Toprak, H. Ü. S. E. Y. İ. N., Yucel, A. Y. T. A. Ç., Begec, Z. E. K. İ. N. E., & Ersoy, O. (2007). Control of shivering during regional anaesthesia: prophylactic ketamine and granisetron. *Acta anaesthesiologica scandinavica*, 51(1), 44-49.
8. Misiran, K., & Aziz, F. Z. (2013). Effectiveness of low-dose midazolam plus ketamine in the prevention of shivering during spinal anaesthesia for emergency lower limb surgery. *Southern African Journal of Anaesthesia and Analgesia*, 19(3), 164-170.
9. Dal, D., Kose, A., Honca, M., Akinci, S. B., Basgul, E., & Aypar, U. (2005). Efficacy of prophylactic ketamine in preventing postoperative shivering. *British journal of anaesthesia*, 95(2), 189-192.
10. Cattaneo, C. G., Frank, S. M., Hesel, T. W., El-Rahmany, H. K., Kim, L. J., & Tran, K. M. (2000). The accuracy and precision of body temperature monitoring methods during regional and general anesthesia. *Anesthesia & Analgesia*, 90(4), 938-945.
11. Tsai, Y. C., & Chu, K. S. (2001). A comparison of tramadol, amitriptyline, and meperidine for postepidural anesthetic shivering in parturients. *Anesthesia & Analgesia*, 93(5), 1288-1292.
12. Sagir, O., Gulhas, N. U. R. Ç. İ. N., Toprak, H. Ü. S. E. Y. İ. N., Yucel, A. Y. T. A. Ç., Begec, Z. E. K. İ. N. E., & Ersoy, O. (2007). Control of shivering during regional anaesthesia: prophylactic ketamine and granisetron. *Acta anaesthesiologica scandinavica*, 51(1), 44-49.
13. Gangopadhyay, S., Gupta, K., Acharjee, S., Nayak, S. K., Dawn, S., & Piplai, G. (2010). Ketamine, tramadol and pethidine in prophylaxis of shivering during spinal anaesthesia. *Journal of Anaesthesiology Clinical Pharmacology*, 26(1), 59-63.
14. Kose, E. A., Honca, M., Dal, D., Akinci, S. B., & Aypar, U. (2013). Prophylactic ketamine to prevent shivering in parturients undergoing Cesarean delivery during spinal anesthesia. *Journal of clinical anesthesia*, 25(4), 275-280.
15. Pazuki SH. Noruzi A. shademan A (2011) Compare of effective dose of ketamine and pethedin for decrease shivering post operation.

- Scientific journal of Arak university .YEAR12.N47 9-6.
16. Katzong Bertram J, Bemasters Suzanov. Translator: Rezvafard Mehrnaz, Sinai Farnaz. Basic and Clinical Pharmacology (2009). first volume. First Edition, 2010. Negaresh Pres.
  17. Sigel, E., & Steinmann, M. E. (2012). Structure, function, and modulation of GABAA receptors. *Journal of Biological Chemistry*, 287(48), 40224-40231.
  18. Kanto, J. H. (1985). Midazolam: the first water-soluble benzodiazepine; pharmacology, pharmacokinetics and efficacy in insomnia and anesthesia. *Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy*, 5(3), 138-155.
  19. Dundee, J. W., Halliday, N. J., Harper, K. W., & Brogden, R. N. (1984). Midazolam: a review of its pharmacological properties and therapeutic use. *Drugs*, 28, 519-543.
  20. Vranken, J. H., Troost, D., Wegener, J. T., Kruis, M. R., & Van der Vegt, M. H. (2005). Neuropathological findings after continuous intrathecal administration of S (+)-ketamine for the management of neuropathic cancer pain. *Pain*, 117(1-2), 231-235.
  21. Fitzgibbons RJ, Forse, RA .Clinical practice. Groin hernias in adults". *The New England Journal of Medicine*.2015. 372 (8): 756–63.
  22. PACKARD, G. B. (1963). Inguinal hernia of infancy and childhood. *Archives of Surgery*, 86(2), 299-303.
  23. Kamtoh, G., Pach, R., Kibil, W., Matyja, A., Solecki, R., Banas, B., & Kulig, J. (2014). Effectiveness of mesh hernioplasty in incarcerated inguinal hernias. *Videosurgery and Other Miniinvasive Techniques*, 9(3), 415-419.
  24. Manuel C, Paradou. Ronaldo Dee Miller. Principles of Miller Anesthesia. Translator: Dr. Farhad Etezad and Dr. Elham Fakharzadeh Naeini. First edition, 1397, Arjmand Publishing.
  25. Okhovat pour H. A review of pharmacology, University of Tehran, 1396. Lotus publications, (Persian).
  26. Bahrami G, Aram SH, Jabal ameli M. The effect of lidocaine and bupivacaine on abdominal cavity on severity of pain and symptoms of patients after abdominal hysterectomy. *Journal of Mazandaran University of Medical Sciences*, 2007; 15(53):1-8.(Persian).
  27. Reves, J. D., Fragen, R. J., Vinik, H. R., & Greenblatt, D. J. (1985). Midazolam: pharmacology and uses. *Anesthesiology*, 62(3), 310-324.
  28. D. M. Little Jr., *Classical Anaesthesia Files*, Wood Library—Museum of Anesthesiology, 1985. 30A