

# The preventive and predictive value of simple exercises (Stair-climbing, squats, and push-ups): An educational article and expert opinion

Aamir Jalal Al-Mosawi

Advisor in Pediatrics and Pediatric Psychiatry, Children Teaching Hospital of Baghdad Medical City and the National Training and Development Center.

**Corresponding Author:** Aamir Jalal Al-Mosawi, Advisor in Pediatrics and Pediatric Psychiatry, Children Teaching Hospital of Baghdad Medical City and the National Training and Development Center.

**Received Date:** 31 December 2022; **Accepted Date:** 06 January 2023; **Published date:** 16 January 2023

**Citation:** Aamir Jalal Al-Mosawi, (2023). The preventive and predictive value of simple exercises (Stair-climbing, squat, and push-ups): An educational article and expert opinion. Journal of Obesity and Fitness Management. 2(1). DOI: 10.58489/2836-5070/004

**Copyright:** © 2023 Aamir Jalal Al-Mosawi, this is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

Cardiovascular diseases including ischemic heart diseases continue to be the leading cause of mortality in Iraq and many other countries in the world.

As early as the 1980s, the relationship between increased physical fitness and reduction of morbidity and mortality from chronic disease, particularly cardiovascular diseases have been increasingly recognized. Therefore, the need for simple preventive exercises and predictive exercise tests has been increasingly demanded.

The aim of this educational mini-review is to highlight the preventive and predictive values of simple exercises (Stair-climbing, squat, and push-up), and also to present the relevant expert opinion.

**Keywords:** Physical fitness, cardiovascular diseases, expert opinion.

## Introduction

Cardiovascular diseases including ischemic heart diseases continue to be the leading cause of mortality in Iraq and many other countries in the world [1,2,3].

As early as 1984, Bruce emphasized the relationship between increased physical fitness and reduction of morbidity and mortality from chronic disease, particularly cardiovascular diseases [4]. Therefore, the need for simple preventive and predictive exercise has been increasingly demanded.

As early as 2002, Kong Chuan Teh (Figure-1A) and Abdul Rashid Aziz from Singapore reported a study that suggested a simple stair-climbing exercise is a feasible exercise for encouraging physical activity for its cardiopulmonary benefits [5].

In 2005, Colin Boreham (Figure-1B) from Northern Ireland and his research group reported an eight-week controlled study which included 15 females with mean age of 18.8 years. The study showed that several short sessions of stair climbing during the day can help in reducing cardiovascular risk factors [6].



**Fig-1A:** Kong Chuan Teh from Singapore



**Fig 1B:** Colin Boreham from Northern Ireland

## Journal of Obesity and Fitness Management

In 2018, Yi Guo (Figure-1C) and his research group from the United States reported a cross-sectional study which included forty individuals (18 males and 22 females) aged 18 to 64 years (Mean age: 31.2 years). The participants performed Ruffier squat test and the Balke maximal treadmill test. The study showed that the squat exercise is a valid clinical test for the evaluation of cardiopulmonary fitness in a primary care setting [7].



**Fig 1C:** Yi Guo from the United States

In 2019, Justin Yang (Figure-1D) from Boston, Massachusetts and his international research group from the United States, Cyprus and Italy reported a retrospective study that included 1104 male firefighters (Aged 18 years or older) observed during the period from the 1st of January, 2000 to the 31st of December 2010, and had push-up capacity data. The participants were followed for up to ten years and the results of the final analysis were available on the 11th of August, 2018. The study found an important negative association between higher baseline push-up capacity and cardiovascular disease events. Participants who were unable to perform ten push-ups had the greatest risk of experiencing cardiovascular disease events [8].



**Fig 1D:** Justin Yang from Boston, Massachusetts

In 2021, Anna C Whittaker (Figure-1E) and her research group from the United Kingdom emphasized the available evidence suggesting that stair climbing exercise and physical activity can be associated with a better lipoprotein profile, a healthier blood pressure, lower body weight, and a higher aerobic physical fitness. They reported a study which suggested that that daily stair climbing can be protect against the

metabolic syndrome which is associated with unhealthy lipid profiles, blood sugar levels, blood pressure and obesity [9].



**Fig 1E:** Anna C Whittaker from the United Kingdom

The **Ruffier Squat** test which was developed by James-Edward Ruffier (Figure-2) during the early 1950s to test the cardiovascular fitness of athletes, and its modified version suggested by Dickson are not particularly suitable for use by physicians caring for non-athletic individuals.

**The Ruffier Squat** test involves recording heart rate before and after performing 30 squats in 45 seconds. Most untrained individuals visiting clinician can never achieve the capacity of 30 squats /45 seconds.



**Fig 2:** James Edward Ruffier (1875-1965), a French physician  
Expert opinion

Based on the available evidence and scientific ideas, it is recommended that clinicians keep records of the baseline of simple exercises (Stair-climbing, squat, and push-up) capacity of the individuals under their care to help in the future identification of increased risk of cardiovascular disease and also help in designing an appropriate preventive program.

It is also recommended to encourage people to maintain a good capacity of these simple exercises, which in our opinion include 4 times ascending/descending 15 steps-stair in one minute, 25-30 squats in one minute, and 20-25 push-ups in one minute.

### Acknowledgment

The author has the copyright of all the sketches included in this paper.

## References

1. Al-Mosawi AJ. Healthcare system in Iraq .1st ed., Saarbrücken; LAP Lambert Academic Publishing: 2014 (ISBN: 978-3-659-55533-6).
2. Al-Mosawi AJ. Medical and Healthcare leadership: A training course. 1st ed., Saarbrücken; LAP Lambert Academic Publishing: 2016 (ISBN: 978-3-659-85373-9).
3. Al-Mosawi, A. J. (2021). Mortality pattern in Iraq before covid-19 pandemic. *Community Medicine and Health Education Research*, 2(1), 11-16.
4. Bruce, R. A. (1984). Exercise, functional aerobic capacity, and aging--another viewpoint. *Medicine and Science in Sports and Exercise*, 16(1), 8-13.
5. Teh, K. C., & Aziz, A. R. (2002). Heart rate, oxygen uptake, and energy cost of ascending and descending the stairs. *Medicine & Science in Sports & Exercise*, 34(4), 695-699.
6. Boreham CA, Kennedy RA, Murphy MH, Tully M, Wallace WF, Young I. Training effects of short bouts of stair climbing on cardiorespiratory fitness, blood lipids, and homocysteine in sedentary young women. *Br J Sports Med* 2005 Sep;39(9):590-3. Doi: 10.1136/bjism.2002.001131.
7. Guo Y, Bian J, Li Q, Leavitt T, Rosenberg EI, Buford TW, Smith MD, Vincent HK, Modave F. A 3-minute test of cardiorespiratory fitness for use in primary care clinics. *PLoS One* 2018 Jul 30;13(7): e0201598. Doi: 10.1371/journal.pone.0201598.
8. Yang J, Christophi CA, Farioli A, Baur DM, Moffatt S, Zollinger TW, Kales SN. Association Between Push-up Exercise Capacity and Future Cardiovascular Events Among Active Adult Men. *JAMA Netw Open* 2019 Feb 1;2(2): e188341. Doi: 10.1001/jamanetworkopen.2018.8341.
9. Whittaker, A. C., Eves, F. F., Carroll, D., Roseboom, T. J., Ginty, A. T., Painter, R. C., & de Rooij, S. R. (2021). Daily stair climbing is associated with decreased risk for the metabolic syndrome. *BMC Public Health*, 21(1), 923.
10. Dickson J. the use of the cardiac index of Ruffier in the medical control of sportsmen [in French]. *Med Educ Phys Sport* 1950; 2:65–79.7.
11. Ruffier JE. Considerations on the index of heart resistance to effort [in French]. *Med Educ Phys Sport*. 1951; 3:7-12.