

# Study of vinpocetine adsorption on the surface of liposomes obtained from soya lecithine

Yu. A. Polkovnikova

Federal State Budget Educational Institution of Higher Education "Voronezh State University," Russia, 394036, Voronezh, Russia

**\*Corresponding Author:** Yu. A. Polkovnikova, Federal State Budget Educational Institution of Higher Education "Voronezh"

**Received:** 30 November 2022; **Accepted:** 12 December 2022; **Published:** 22 December 2022.

**Citation:** Yu. A. Polkovnikova, (2022). Study of vinpocetine adsorption on the surface of liposomes obtained from soya lecithine. Journal of Pharmacy and Drug Development. 1(3). DOI: 10.58489/2836-2322/012

**Copyright:** © 2022 Yu. A. Polkovnikova, this is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

Liposomes are one of the best studied types of nanoparticles that are considered as contemporary and efficient means for delivery of various medicinal agents widely applied in the clinical practice. Taking into account the features of transportation, translocation through histological barriers, cellular membranes and metabolic transformations liposomal pharmaceutical drugs show unique properties, first of all connected with the distinctions in their pharmacokinetics. Determination of adsorption characteristics for vinpocetine applied on the surface of liposomes obtained from soya lecithine. Liposomes from soya lecithine were obtained by hydration/rehydration technique. In order to study characteristics of vinpocetine adsorption on the surface of liposomes technique of equilibrium dialysis was utilized. Sol of iron hydroxide was used as a comparison adsorbent. Separation of the dispersive medium from sol colloid particles was performed with the use of column chromatography. Comparative investigations of vinpocetine adsorption were performed both on comparison adsorbent and on the liposomes obtained from soya lecithine. Rather efficient adsorption of vinpocetine on the liposomes at their low concentrations was observed. Vinpocetine adsorption, liposome, soya lecithine

**Keywords:** neurons; cerebral ischemia; L-arginine; Omega-3 PUFAs

## Introduction

Nanocarriers used for the drug delivery to the organs and tissues differ by their size, shape and composite materials [1]. The properties of each nanoparticle are defined by the extent of its payload with a drug, stability, the rate of the drug release and the presence of a ligand for the directed transportation [2,3].

In order to get the required therapeutic effect pharmaceutical substance encapsulated into the vesicles should be available to the targeted cells [4]. At this point liposomes differ from the other controlled delivery systems where the biologically active compounds are released either in the blood plasma or just in the place of injection. Then the captured preparation can be selectively accumulated within the affected part of an organism due to the passive (unintended) or active targeting [5].

Liposomes are spherical vesicle structures composed of a uni- or multilamellar lipid bilayer surrounding internal aqueous compartments and a

relatively impermeable outer lipophilic phospholipid bilayer. Liposomes have gained considerable attention as drug delivery carriers because they are biocompatible, nontoxic, can deliver both hydrophilic and lipophilic drug molecules, protect their cargo from degradation by plasma enzymes, and transport their load across biological membranes and the blood brain barrier [6].

When using liposomes as an independent preparation or as a carrier for medicinal agents it is required to study the effect of liposomes on the cells in human organism. The determining factor in this case can be as the chemical composition of liposomes as their size [7].

Vinpocetine is a vasoactive and nootropic preparation that proves to be semisynthetic derivative of the common periwinkle plant alkaloid. Vinpocetine is referred to the type of compounds that are practically insoluble in water [8].

Objective of the investigation: determination of

## Pharmacy and Drug Development

characteristics for adsorption of vincopetine on the surface of liposomes obtained from soya lecithine.


### Materials and Methods

#### **The study of vincopetine adsorption characteristics on the surface of liposomes**

Characteristics of vincopetine adsorption on the surface of liposomes were studied by the equilibrium dialysis technique [9]. The choice of this method is stipulated by the fact that quantitative analysis of the equilibrium vincopetine concentration in the dispersive medium necessary for the determination of the adsorption value is complicated by the presence

of dispersive phase - liposomes. Semipermeable membrane with diameter of pores that is sufficient for the penetration of vincopetine molecules but leakless for liposomes ensures obtaining of vincopetine solution with the concentration close to that one vincopetine in the liposomes dispersive medium. Solution prepared in such a way can be analyzed quantitatively with the use of spectrophotometry.

In order to determine the linearity of the applied photometric analysis technique a calibration graph was plotted within the range of concentrations and optical densities required for the performed study at the wavelength of 313 nm (figure 1).



**Fig 1:** Calibration plot for the quantitative determination of vincopetine in the medium of 0,01 M of aqueous solution of the hydrochloric acid by spectrophotometry technique at the wavelength of 313 nm

In this study the values of vincopetine adsorption on liposomes at the different concentrations of vincopetine were determined as well as the constants in Freundlich and Langmuir equations.

To solve this problem colloid solution of liposomes from soya lecithine was prepared and then its mass fraction was determined. After that the process standard sample (PSS) of vincopetine was prepared and calibration graph was plotted and molar absorption coefficient of vincopetine was determined. Optimal technique of equilibrium dialysis was chosen, equilibrium dialysis for different concentrations of vincopetine was performed and analysis of the obtained results was executed.

Preparation of the liposomes samples prepared from soya lecithine

To obtain liposomes from soya lecithine hydration/rehydration was applied. Soya lecithine solution (Sigma) in ethyl alcohol was evaporated in the rotor

evaporator at the temperature of 45°C and pressure of -0,085 MPa. Then 0,01 M solution of hydrochloric acid (pH=2,0) was added. For obtaining of liposomes solutions were subjected to irradiation with ultrasound disintegrator for 15 minutes. Next, liposomes were filtered through glass filter with diameter of pores 16 m.

#### **Preparation of the process standard sample (PSS) of vincopetine**

An accurate sample of 12 mg of vincopetine was solved in 0,01 M aqueous solution of hydrochloric acid in the graduated flask of 50 ml volume and was brought up to the label with a solvent.

Equilibrium dialysis was performed with the use of dialysis vials Easy Dial-L having semipermeable membrane and its passing characteristic was of 14 kDa. For the determination of vincopetine adsorption value by liposomes the main experiment was made (dialyzer A), where adsorption was observed,

reference experiment (dialyzer B) where dialysis took place but there was no liposomes, and the experiment for the measurement of content of the free lecithine in the dispersive medium (dialyzer C). 12 ml of liposome solution and dialysis vial were placed into dialyzer A. Note, that 3 ml of vinpocetine solution with the concentration  $C_0$  was placed into the dialyzer vial. 12 ml of 0,01 M solution of hydrochloric acid and dialysis vial were placed inside dialyzer B. After that 3 ml of vinpocetine solution with the concentration of  $C_0$  was placed into dialyzer vial. And 12 ml of liposome solution and the dialyzer vial were placed into dialyzer C, while in the dialyzer vial 3 ml of 0,01 M solution of hydrochloric acid was placed as well. Dialysis was performed in thermostat at the temperature of 37 °C for 12 hours. After that using spectrophotometer LOMO SF-56 optical density of solutions was measured in the dialyzer vials previously placed in the dialyzers A, B and C at the wavelength of 313 nm (DA, DB and DC, respectively). For the determination of vinpocetine concentration in the dialyzer A the difference of optical densities  $D = DA - DB$  was applied.

## Results

### *Study of the adsorption characteristics of vinpocetine on the surface of liposomes*

Taking into account vinpocetine solubility in the acidic medium as a dispersive medium for liposomes 0,01 M solution of hydrochloric acid was chosen as such a medium. Vinpocetine is also soluble in ethanol, but during liposome interaction with ethanol their coagulation occurs.

Order of plotting calibration graph and determination of absorptivity for vinpocetine in 0,01 M medium of aqueous solution of hydrochloric acid

During the study of adsorption characteristics of vinpocetine on liposomes the quantitative determination of vinpocetine in 0,01 M aqueous solution of hydrochloric acid was performed by spectrophotometry technique. To do so vinpocetine spectrophotometry spectrum was measured for 0,01 M of vinpocetine in the aqueous solution of hydrochloric acid.

For 0,01 M of aqueous solution of hydrochloric acid ultraviolet spectrum of vinpocetine is characterized by the peaks of optical density at the wavelengths of 268 and 313 nm. The choice of the maximum in the optical density used for the quantitative determination of vinpocetine was made basing on the possible influence of the free lecithine used as a dispersive

medium for liposomes capable to penetrate through semipermeable membrane [20]. To check this idea a test dialysis was performed where 3 ml of solution with liposomes were placed into dialysis vial while 12 ml of 0,01 M of the aqueous solution with hydrochloric acid were placed into a dialyzer. Dialysis was performed for 12 hours at 37° C. Then ultraviolet absorption spectrum of solution in dialyzer was measured. For the analysis of the quantitative content of vinpocetine the peak corresponding to 313 nm maximum of absorption was chosen since in this spectral range the contribution of lecithine in the dispersive medium for liposomes in the optical density proves to be minimal.

In order to determine the linearity of the applied photometric analysis technique a calibration graph was plotted within the range of concentrations and optical densities required for the performed study at the wavelength of 313 nm.

The choice of technique for the equilibrium dialysis in order to study the process of adsorption for vinpocetine on liposomes

The objective of the choice of technique for the equilibrium dialysis [20] was the search for conditions when the change of vinpocetine concentration due to its adsorption on liposomes would be maximum one. In the process of the choice of technique concentrations of vinpocetine in a dialyzer varied as well as the volumes of liquids by both sides of the dialyzer membrane.

To make the study of the effect of vinpocetine concentration on the process of its adsorption on the liposomes 7 dialyzers were prepared. In the dialyzers 1 – 5 were OSS solutions were placed according to the Table 6, with a volume of 12 ml. 12 ml of solution 5 from the Table 6 was placed into dialyzer 6. 12 ml of the aqueous solution of hydrochloric acid with a concentration of 0.01 M was placed in a dialyzer 7. Next, in all of the dialyzers dialysis vials of Easy Dial-L type with semi-permeable membranes and transmission characteristic of 14 kDa were placed. 2 ml of liposome solution were placed into dialysis vials of the dialyzers 1 – 5 and 7. And 2 ml of 0,01 M of the aqueous solution of hydrochloric acid was placed in dialyzer 6. Dialysis was performed in a thermostat at 37°C for 12 hours. Then, the optical density of solutions in the dialyzers was measured. Solutions in the dialyzers 3 – 6 were diluted with 0.01 M solution of the hydrochloric acid before measurements. Results of the measurements are presented in Table

**Table 1:** Results of the equilibrium dialysis for vinpocetine solutions with different concentration in the presence of liposomes

No of dialyzer	Dilution during measurements of the optical density	Optical density at the wavelength of 313 nm	Optical density at the wavelength of 313 nm with the account of dilution and free lecithine	Optical density at the wavelength of 313 nm with the account of dilution for PSS of vinpocetine	Change of the optical density for vinpocetine as a result of adsorption	The presence of coagulation in the dialysis vial
1	1	0,2353	0,2178	0,2582	0,0404	Absence of coagulation
2	1	0,4645	0,4470	0,5164	0,0694	Little flaky sediment
3	5,22	0,1910	0,9795	1,0328	0,0532	Little flaky sediment
4	5,21	0,3653	1,8857	2,0655	0,1798	Almost complete coagulation, solution is close to transparent one
5	10	0,4024	4,0065	4,1310	0,1245	Complete coagulation, transparent solution
6	10	0,4131	4,1310	–	–	–
7	1	0,0175	0,0000	–	–	Absent

Conditional optical density was calculated with the account of dilution of the solutions just before the spectrophotometry measurements. Optical density of solution from the dialyzer 7 was subtracted from the total optical density (implying optical density of the free lecithine).

According to the results of the executed dialysis it is possible to make a conclusion that concentration of vinpocetine with the optical density of solution more than 0,25 and this can result in coagulations of liposomes. If concentration of vinpocetine corresponds to the optical density of 0,25 then the change of the optical density under adsorption on liposomes was of 0,04.

In order to enhance the amount of the adsorbed vinpocetine the volume occupied by liposomes was increased up to 12 ml. And solution of liposomes was placed directly in a dialyzer while 2 ml of vinpocetine was placed into the dialysis vial (dialyzer A). Vinpocetine solution for this experiment was prepared by mixing of OSS solution of vinpocetine and 0,01 M of the aqueous solution of hydrochloric acid with a ratio of 1:1 (solution A). 12 ml of the aqueous solution of 0,01 M of hydrochloric acid was placed into the second dialyzer (dialyzer B) as well as a dialysis vial. 2 ml of solution A was measured and placed into the dialysis vial. 12 ml of liposomes solution was measured and placed into third dialyzer

(dialyzer B) and then placed there dialysis vial with was filled with 2 ml of aqueous solution of 0,01 M of hydrochloric acid. The dialysis was performed in thermostat at the temperature of 37 °C for 12 hours. No coagulation of liposomes was observed in the dialyzers A and B after 12 hours of the experiment.

According to the results of experiment with an increased volume of liposomes solution a technique was chosen for the further investigation which includes placing of 12 ml of the liposomes solution in the dialyzer. After that in order to study adsorption of vinpocetine on liposomes the authors applied vinpocetine concentrations that did not cause liposomes coagulation

Determination of adsorption value for vinpocetine on the liposomes by equilibrium dialysis.

In order to determine adsorption parameters of vinpocetine on the liposomes a series of vinpocetine solutions with different concentrations was prepared for the study. Their dialysis was made according to the technique presented above. Equilibrium concentrations of vinpocetine in the dialyzers A and B were calculated with the use of the molar absorption coefficient.

Next, a calculation of the value of vinpocetine adsorption value on the liposomes was performed in accordance with the results of determination for the equilibrium concentration in the dialyzer vials of the

main and reference experiments.

Under increase if vinpocetine concentration the value of adsorption upraises as well and it attains maximum at the concentrations of about 0,030–0,035 mole/kg.

Results of determination of the characteristics for vinpocetine adsorption on the liposomes are presented in Table 2.

**Table 2:** Results of determination for adsorption characteristics of vinpocetine adsorption on liposomes

Equation	Constant	Value
Freundlich equation	$1/n$	$0,505919 \pm 0,108038364$
	$k$ , mole/kg	$3,615507457 \pm 1,163161619$
Langmuir equation	$A^\infty$ , mole/kg	$0,0122987 \pm 0,005485$
	$b$ , mole/l	$4,08717 \cdot 10^{-6} \pm 6,74916 \cdot 10^{-6}$

### Conclusion

1. Ultimate adsorption of vinpocetine on liposomes is considerably less than that one on the colloid particles of iron hydroxide (III) sol.
2. Constant  $b$  in Langmuir equation (the value of concentration when a half of the ultimate adsorption is attained) proves to be less as compared with that one characteristic for adsorption on the colloid particles of iron hydroxide (III) sol; it means quite efficient adsorption of vinpocetine by liposomes at low concentrations.
3. According to the results of performed investigations one can make a conclusion that incorporation of vinpocetine into liposomes by its addition to the ready-made colloid solution of liposomes is rather low-efficient.

### References

1. W. Brenner, F. Gschnait, A. Hautarzt, 29(7), 392-4 (1978).
2. Polkovnikova, Y. A., Glushko, A. A., Mikhailovskaya, I. Y., & Karieva, Y. S. (2017). Simulation of vinpocetine release process from microcapsules with hydrophobic shell. *Pharmacy & Pharmacology*, 5(4), 344-367.
3. Scholtz, A. W., Hahn, A., Stefflova, B., Medzhidieva, D., Ryazantsev, S. V., Paschinin, A., ... & Weisshaar, G. (2019). Efficacy and safety of a fixed combination of cinnarizine 20 mg and dimenhydrinate 40 mg vs betahistine dihydrochloride 16 mg in patients with peripheral vestibular vertigo: a prospective, multinational, multicenter, double-blind, randomized, non-inferiority clinical trial. *Clinical drug investigation*, 39, 1045-1056.
4. S. Sethi, B. Mangla, S. Kamboj, V.A. Rana Int J Biol Macromol. 1(117), 350-36113 (2018).
5. Shahba, A. A. W., Ahmed, A. R., Alanazi, F. K., Mohsin, K., & Abdel-Rahman, S. I. (2018). Multi-layer self-nanoemulsifying pellets: an innovative drug delivery system for the poorly water-soluble drug cinnarizine. *Aaps Pharmscitech*, 19, 2087-2102.
6. Ammar, H. O., Ghorab, M., Kamel, R., & Salama, A. H. (2016). A trial for the design and optimization of pH-sensitive microparticles for intestinal delivery of cinnarizine. *Drug delivery and translational research*, 6, 195-209.
7. Hou, G., Niu, J., Song, F., Liu, Z., & Liu, S. (2013). Studies on the interactions between ginsenosides and liposome by equilibrium dialysis combined with ultrahigh performance liquid chromatography-tandem mass spectrometry. *Journal of Chromatography B*, 923, 1-7.
8. Dwiastuti, R., Radifar, M., Marchaban, M., Noegrohati, S., & Istyastono, E. P. (2016). Molecular dynamics simulations and empirical observations on soy lecithin liposome preparation. *Indonesian Journal of Chemistry*, 16(2), 222-228.
9. Yu. A. Polkovnikova, A. I. Slivkin, *Pharmaceutical Chemistry Journal*, 50(8), 553-555 (2016).
10. Y.A. Polkovnikova, A.A. Glushko, *Pharmacy & Pharmacology*, 6(2), 197-210 (2018).